Why Toxicity Values Are in Flux: A Summary and Perspective on Recent IRIS Assessments

Anita K. Meyer DABT

Environmental & Munitions CX
Huntsville Engineering and Support Center

E²S² Symposium and Exhibition New Orleans, LA

May 10, 2011









maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding an DMB control number.	ion of information. Send comment arters Services, Directorate for Inf	s regarding this burden estimate formation Operations and Reports	or any other aspect of the property of the pro	his collection of information, Highway, Suite 1204, Arlington
1. REPORT DATE 10 MAY 2011	2. REPORT TYPE			3. DATES COVERED 00-00-2011 to 00-00-2011	
4. TITLE AND SUBTITLE				5a. CONTRACT	NUMBER
Why Toxicity Values Are in Flux: A Summary and Perspective on Rece			ective on Recent	5b. GRANT NUMBER	
IRIS Assessments				5c. PROGRAM F	ELEMENT NUMBER
6. AUTHOR(S)				5d. PROJECT NU	UMBER
				5e. TASK NUMBER	
				5f. WORK UNIT	NUMBER
U.S. Army Corps of	ZATION NAME(S) AND AE of Engineers (USAC) e (EM CX),4820 Un AL,35816	E),Environmental	and Munitions	8. PERFORMING REPORT NUMB	G ORGANIZATION ER
9. SPONSORING/MONITO	RING AGENCY NAME(S) A	AND ADDRESS(ES)		10. SPONSOR/M	IONITOR'S ACRONYM(S)
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAII Approved for publ	LABILITY STATEMENT ic release; distributi	ion unlimited			
	OTES DIA Environment, l 1 in New Orleans, L		Sustainability (E2	S2) Symposi	um & Exhibition
14. ABSTRACT					
15. SUBJECT TERMS					
			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	17	RESI ONSIBLE I ERSON

Report Documentation Page

Form Approved OMB No. 0704-0188

Role of Risk Assessment on Environmental Projects

- Used to determine whether site requires further study or remediation
- Used with regulatory values to determine cleanup levels
- Risk-based screening levels (aka PRGs, RBCs, RSLs)
 - ► Screen sites early in project lifecycle
 - ▶ Determine project quantitation limits



Risk-Based Values Function of Toxicity and Exposure

Risk = Intake Toxicity

- Tiered approach used to identify toxicity values for site risk assessments
- Integrated Risk Information System (IRIS)
- IRIS values also inform regulatory decisions (MCLs etc.)





Emerging Contaminants (ECs)


- Are chemicals or materials of interest that are characterized by:
 - ▶ a perceived or real threat to human health or environment, and
 - ► there is no currently published health standard or there is an existing health standard, but the standard is evolving or being re-evaluated.

Source: "Initiation of Emerging Contaminants Characterization and Response Actions for Protection of Human Health" Issue Paper (ECOS &DoD Sustainability Workgroup, 2008)



DoD Participation in IRIS Interagency Reviews

Assessment Development Process for New IRIS



Potential Changes in Risk Assessment of PAHs

- Relative potency factors used to assess carcinogenicity of PAHs – all set relative to benzo(a)pyrene
- EPA: "Development of a Relative Potency Factor (RPF) Approach for Polycyclic Aromatic Hydrocarbon (PAH) Mixtures"
 - ► Under review by EPA Science Advisory Board
- RPF approach retained but updated by new data/science



Carcinogenic PAHs and Relative Potency Factors

	Current RPF	Draft RPF
Benzo(a)pyrene	1	1
Benz(a)anthracene	0.1	0.2
Benzo(b)fluoranthene	0.1	0.5
Benzo(k)Fluoranthene	0.01	0.03
Chrysene	0.001	0.1
Dibenz(a,h)anthracene	1	6
Indeno(1,2,3-c,d)pyrene	0.1	0.07





Additional PAHs from 2010 RPF Assessment

- Anthanthrene
- Benzo[g,h,i]perylene
- Benzo[j]fluoranthene
- Cyclopenta[c,d]pyrene
- Dibenzo[a,e]fluoranthene
- Dibenzo[a,e]pyrene
- Dibenzo[a,h]pyrene
- Dibenzo[a,i]pyrene
- Dibenzo[a,l]pyrene
- Fluoranthene

- Benz[b,c]aceanthrylene
- Benz[e]aceanthrylene
- Benz[j]aceanthrylene (60x)
- Benz[l]aceanthrylene
- Cyclopenta[d,e,f]chrysene
- Naphtho[2,3-e]pyrene









Tetrachloroethylene Status: External Peer Review

- 1998 initiated
- June 2008 external review version released
- Nat'l Academy
 Review Feb 2006
- Current external (SAB) and public review

Risk-Based Screening Levels*			
	Res. Soil (mg/kg)	Res. Water Use (µg/L)	Indoor Air (µg/m³)
Current	0.55	0.11	0.41
New (draft)	0.293	0.179	0.122
Sources of current toxicity values include EPA IRIS, ATSDR and CalEPA. Lowest RSL target risk = 10 ⁻⁶ . Draft values not suitable for project use.			

Trichloroethylene Status: External Peer Review

- IRIS values withdrawn late 1980s
- 2001 draft released
- 2006 NAS review
- 2009 re-released
- Lowest RSLs based on 10⁻⁶ cancer risk

Risk-Based Screening Levels*			
	Res. Soil (mg/kg)	Res. Water Use (µg/L)	Indoor Air (µg/m³)
Current	2.8	2	1.2
New (draft)	0.48	0.23	0.24
CalEPA source of current values. Lowest RSL target risk = 10 ⁻⁶ . Draft			

values not suitable for project use.



Dioxin Status: External Peer Review

- Assessment initiated in 1990
- Released for public and peer review 2010
- Using cancer toxicity values in this draft residential screening levels could be as low as 0.45 ppt dioxin toxicity equivalent (TEQ, for dioxin-like compounds)
- Current EPA policy recommends 1000 ppt
 - ► EPA may release an Interim PRG



Hexavalent Chromium Status: External Peer Review

- Derives an oral slope factor using 2008
 National Toxicology
 Program studies
- NTP studies also used to update noncancer ingestion value (3x change)
- Lowest RSLs based on 10⁻⁶ cancer risk

Risk-Based Screening Levels*			
	Res. Soil (mg/kg)	Res. Water Use (µg/L)	
Old	2300	11	
New (draft)	0.29	0.043	
Draft values not suitable for project use. Cancer risk drives new RSLs.			



Final IRIS Reference Dose for Cis-1,2-Dichloroethylene (September 2010)

- Oral noncancer reference dose (RfD) = 0.002 mg/kg-day; increased kidney weight
 - PPRTV used until this time
 - ► MCL = 70 µg/L
- No cancer toxicity values published

Risk-Based Screening Levels*			
	Residential Soil (mg/kg)	Residential Water Use (µg/L)	
Old	780	370	
New	156	73	
Fold Change		5x	



Final IRIS Reference Dose for Trans-1,2- Dichloroethylene (September 2010)

- Oral noncancer reference dose (RfD) = 0.02 mg/kg-day;
- Based on decreased antibody production by the spleen
- No change in RfD value, but change of critical effect
- MCL = 100 μg/L
- No cancer toxicity values published



Final IRIS Values for 1,4-Dioxane (August 2010)

- Oral noncancer reference dose (RfD) = 0.03 mg/kg-day; Liver & kidney effects
 - ► Inhalation RfC will be developed in separate document
- Oral cancer slope factor (SF) = 0.1 mg/kg-day ⁻¹;
 Rodent liver tumors

Risk-Based Screening Levels*			
	Residential Soil (mg/kg)	Residential Water Use (µg/L)	
Old	44.1	6.1	
New	4.85	0.67	
Fold Change		9x	



Basis for Changes


- New study data
 - ► Sensitive measurements
- Science policy
 - ► Mutagenic mode action adjustment
 - ► Uncertainty factor application
 - ▶ Benchmark Dose
- Other impacts
 - ► Critical study selection
 - ► Linear low dose extrapolation






High-Throughput Screening Assays

batch testing of chemicals for pharmacological/toxicological endpoints using automated liquid handling, detectors, and data acquisition







The future!
EPA Next Generation Risk
Assessment
New technology will lead to greater
capacity and speed. Interpretation of
results is a challenge; studies
underway to link known effects with
screening assay data.

Systems Exposure Science : Extending Network Analysis

Consider coupled networks spanning multiple levels of biological organization

